



# The Uniform over the Whole Line $\mathbb{R}$ Estimates of Spectral Expansions Related to the Selfadjoint Extensions of the Hill Operator and of the Schrödinger Operator with a Bounded and Measurable Potential

I. ANTONIOU

International Solvay Institute for Physics and Chemistry  
ULB, C.P. 231, Boulevard du Triomphe, 1050 Brussels, Belgium  
and

Theoretische Natuurkunde, Free University of Brussels  
C.P. 231, Boulevard du Triomphe, 1050 Brussels, Belgium

V. A. IL'IN

International Solvay Institute for Physics and Chemistry  
ULB, C.P. 231, Boulevard du Triomphe, 1050 Brussels, Belgium  
and

Department of Computational Mathematics and Cybernetics, Moscow State University  
Vorobyovy gory, Moscow 119899, Russia (permanent address)

ilin@cmc.msk.su

**Abstract**—We consider some properties of the spectral expansions related to selfadjoint extensions of the operator  $Hu = -u'' + q(x)u$  over the whole line  $\mathbb{R}$  in the case when  $q(x)$  is a continuous periodic function (the Hill operator) and in the case when  $q(x)$  is a bounded measurable function. This paper gives a brief description of results obtained in the following directions: the uniform over  $\mathbb{R}$  estimates of the generalized eigenfunctions, the uniform over  $\mathbb{R}$  estimates of the spectral function, the uniform over  $\mathbb{R}$  equiconvergence with the Fourier integral expansion, and the uniform over  $\mathbb{R}$  rate of convergence for functions from the Sobolev-Liouville classes.

**Keywords**—Hill operator, One-dimensional Schrödinger operator, Spectral expansion.

The study of scattering on periodic structures demands the necessity for developing the spectral theory of the Hill operator over the whole line  $\mathbb{R} = (-\infty, \infty)$  and, as its generalization, the spectral theory of the Schrödinger operator  $Hu = -u'' + q(x)u$  with an arbitrary bounded and measurable over  $\mathbb{R}$  potential  $q(x)$ .

In this paper, we review our results obtained in this direction. They relate mainly to the following problems:

- (1) the problem of obtaining *the uniform over the whole line  $\mathbb{R}$*  and sharp with respect to the order estimate for the integral of squares of generalized eigenfunctions of the selfadjoint extension of the Hill operator over  $\mathbb{R}$ ;

---

This work was supported by the Commission of the European Communities—DG III/ESPRIT Project—ACTCS 9282. The authors are grateful to L. V. Kritskov for several editorial remarks.

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{T}\mathcal{E}\mathcal{X}$

- (2) the problem of obtaining *the uniform with respect to*  $(x, y) \in \mathbb{R} \times \mathbb{R}$  estimate of the spectral function  $\theta(\lambda; x, y)$  of the selfadjoint extension of the Hill operator over  $\mathbb{R}$ , and the uniform with respect to  $x \in \mathbb{R}$  estimate of the  $L_q(\mathbb{R})$ -norm of  $\theta(\lambda; x, y)$  in variable  $y$  ( $2 \leq q \leq \infty$ );
- (3) the problem of obtaining *the uniform over the whole line*  $\mathbb{R}$  equiconvergence of the spectral expansion of an arbitrary function  $f(x)$  from the class  $L_p(\mathbb{R})$  ( $1 \leq p \leq 2$ ), related to the selfadjoint extension of the Hill operator over  $\mathbb{R}$ , with the conventional Fourier integral expansion of the same function  $f(x)$ ;
- (4) the problem of obtaining for an arbitrary function  $f(x)$  from the Sobolev-Liouville class  $L_2^\alpha(\mathbb{R})$  with  $\alpha \in (1/2, 2]$  *the uniform over the whole line*  $\mathbb{R}$  estimate of deviation of this function  $f(x)$  from its spectral expansion related to the selfadjoint extension of the Hill operator over  $\mathbb{R}$  or related to the Schrödinger operator with a bounded and measurable potential.

Note that there are a lot of papers and reviews which study different aspects of the spectral theory of the Hill operator (see, e.g., [1–7]). But only in [8,9], we managed to estimate the spectral function of the selfadjoint extension of the Hill operator, the related spectral expansion, and the deviation of this spectral expansion from the decomposed function in *the uniform over the whole line*  $\mathbb{R}$  metric.

Let us describe and briefly analyse the obtained results.

It is known (see, e.g., [5, Chapter X.12]) that the Hill operator

$$Hu = -u'' + q(x) \cdot u, \quad (1)$$

with a continuous potential  $q(x)$  satisfying periodic condition  $q(x+a) = q(x)$ , has the unique selfadjoint extension  $\mathcal{L}$  over  $\mathbb{R}$  that is bounded below. Since addition of an arbitrary constant to the potential  $q(x)$  does not alter either its periodicity or the nature of the spectral expansion, but only shifts the spectrum, we assume without loss of generality that extension  $\mathcal{L}$  is bounded below by 1.

It is also known (see [6, Chapter XIII.16]) that the spectral function  $\theta(\lambda; x, y)$  of the selfadjoint extension  $\mathcal{L}$  has the form

$$\theta(\lambda; x, y) = \frac{1}{\pi} \int_{-\infty}^{\lambda} \operatorname{Re} [\psi_t(x) \cdot \overline{\psi_t(y)}] dp(t), \quad (2)$$

where  $p(t)$  is the so-called *quasi-momentum* (it is a continuous nondecreasing function over  $\mathbb{R}$ ), and  $\psi_\lambda(x)$  and  $\overline{\psi_\lambda}(x)$  are *the Bloch eigenfunctions* that for any  $\lambda$  from the spectrum satisfy the equality  $L\psi = \lambda\psi$  and can be obtained for a given  $\lambda$  as eigenfunctions of the monodromy operator  $M(\lambda)$  with eigenvalues  $\exp[\pm i p(\lambda)a]$ .<sup>1</sup>

Furthermore, the spectrum of  $\mathcal{L}$  is absolutely continuous and consists of an infinite number of bands:

$$[\lambda_0, \mu_0], \quad [\lambda_1, \mu_1], \quad [\lambda_2, \mu_2], \dots$$

The sequence  $\lambda_0 \leq \lambda_1 \leq \lambda_2 \leq \dots$  is formed by eigenvalues of the periodic (with period  $a$ ) problem for the Hill operator (1), and the sequence  $\mu_0 \leq \mu_1 \leq \mu_2 \leq \dots$  is formed by eigenvalues of the antiperiodic (with the same period  $a$ ) problem for (1). The qualitative characterization of the quasimomentum  $p(\lambda)$  is as follows: it equals zero on the initial gap  $(-\infty, \lambda_0)$ , equals  $l\pi/a$

<sup>1</sup>The monodromy operator  $M(\lambda)$  for a given  $\lambda$  is an operator which has (in the basis of solutions  $c(x, \lambda)$  and  $s(x, \lambda)$  of  $Lu = \lambda u$ , satisfying the initial conditions  $c(0, \lambda) = s'(0, \lambda) = 1$ ,  $s(0, \lambda) = c'(0, \lambda) = 0$ ) the following form:

$$\begin{pmatrix} c(a, \lambda) & s(a, \lambda) \\ c'(a, \lambda) & s'(a, \lambda) \end{pmatrix}.$$

The trace of this matrix  $D(\lambda) = c(a, \lambda) + s'(a, \lambda)$  is called the Hill discriminant.

on the  $l^{\text{th}}$  gap (the numbering includes the gaps that degenerate into points), and satisfies the equation  $2 \cos[p(\lambda)a] = D(\lambda)$  on the bands (here  $D(\lambda)$  is the Hill discriminant).

Taking into account the Browder-Gårding-Mautner Theorem on the ordered spectral representation of  $L_2(\mathbb{R})$  space with respect to  $\mathcal{L}$  (see, e.g., [10, Chapter XIII.5]) and using formula (2) for the spectral function  $\theta(\lambda; x, y)$  of this extension, we conclude that the quasimomentum  $p(\lambda)$  is the spectral measure of this spectral representation, and functions  $u_1(x, \lambda)$  and  $u_2(x, \lambda)$  which are defined on the bands by the equalities

$$u_1(x, t) = \frac{1}{\sqrt{\pi}} \operatorname{Re} [\psi_t(x)], \quad u_2(x, t) = \frac{1}{\sqrt{\pi}} \operatorname{Im} [\psi_t(x)], \quad (3)$$

and arbitrarily on the gaps,<sup>2</sup> are the kernels (or the generalized eigenfunctions).

If the Fourier transforms of an arbitrary function  $f(x)$  from  $L_p(\mathbb{R})$ ,  $1 \leq p \leq 2$  are defined as

$$\widehat{f}_i(\lambda) = \int_{-\infty}^{\infty} f(y) u_i(y, \lambda) dy, \quad i = 1, 2 \quad (4)$$

(in case  $1 < p \leq 2$ , the integral (4) has meaning in the principal value sense), then the spectral expansion of  $f(x)$  takes the form<sup>3</sup>

$$\sigma_\lambda(x, f) = \sum_{i=1}^2 \int_{\lambda_0}^{\lambda} \widehat{f}_i(t) u_i(x, t) dp(t). \quad (5)$$

The Browder-Gårding-Mautner Theorem states that the spectral expansion (5) of an arbitrary function  $f(x)$  in  $L_2(\mathbb{R})$  converges to  $f(x)$  in  $L_2(\mathbb{R})$  metric as  $\lambda \rightarrow \infty$ , and the Parseval equality holds:

$$\sum_{i=1}^2 \int_{\lambda_0}^{\infty} \widehat{f}_i^2(t) dp(t) = \int_{-\infty}^{\infty} f^2(x) dx. \quad (6)$$

The following statement forms the main result of [8].

**MAIN THEOREM 1.** *Let  $p \in [1; 2]$  be fixed. If  $f(x)$  is an arbitrary function from  $L_p(\mathbb{R})$ ,  $\sigma_\lambda(x, f)$  is the spectral expansion of  $f(x)$  related to the selfadjoint extension  $\mathcal{L}$  of the Hill operator (1) over  $\mathbb{R}$ ,  $S_\lambda(x, f)$  is the Fourier integral expansion of  $f(x)$ , namely*

$$S_\lambda(x, f) = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{\sin [\sqrt{\lambda}(x-y)]}{x-y} f(y) dy, \quad (7)$$

then

$$\lim_{\lambda \rightarrow \infty} |\sigma_\lambda(x, f) - S_\lambda(x, f)| = 0,$$

uniformly with respect to  $x$  over the whole line  $\mathbb{R}$ .

This theorem yields that any convergence criterion of the Fourier integral expansion for a function in  $L_p(\mathbb{R})$ ,  $1 \leq p \leq 2$ , at a point or on a set can serve as a convergence criterion (at a point or, respectively, on a set) of the spectral expansion related to the selfadjoint extension  $\mathcal{L}$  of the Hill operator (1) over  $\mathbb{R}$ .

The proof of Main Theorem 1 essentially uses two auxiliary assertions which have a separate mathematical interest.

<sup>2</sup>Functions  $u_1(x, \lambda)$  and  $u_2(x, \lambda)$  can be defined arbitrarily on the gaps because there we have  $dp(\lambda) \equiv 0$ . These functions only must satisfy  $Lu_1(x, \lambda) = \lambda u_1(x, \lambda)$  and  $Lu_2(x, \lambda) = \lambda u_2(x, \lambda)$ .

<sup>3</sup>We take  $\lambda_0$  instead of  $-\infty$  as the lower bound in (5) because everywhere on the gap  $(-\infty, \lambda_0) : dp(t) \equiv 0$ . Note also that  $\lambda_0 \geq 1$ .

ASSERTION 1. If  $u_1(x, \lambda)$  and  $u_2(x, \lambda)$  are the generalized eigenfunctions (3) of the ordered spectral representation of  $L_2(\mathbb{R})$  with respect to the selfadjoint extension  $\mathcal{L}$  of the Hill operator (1) over  $\mathbb{R}$ , then for all  $\tau \geq \sqrt{\lambda_0} \geq 1$ , the following estimate:

$$\sum_{i=1}^2 \int_{\tau \leq \sqrt{t} \leq \tau+1} |u_i(x, t)|^2 dp(t) = O(1), \quad (8)$$

holds uniformly with respect to  $x \in \mathbb{R}$ .

ASSERTION 2. If  $q$  is an arbitrary fixed number satisfying  $2 \leq q \leq \infty$ ,  $\theta(\lambda; x, y)$  is the spectral function of the selfadjoint extension  $\mathcal{L}$  of the Hill operator (1) over  $\mathbb{R}$ ,

$$\theta_0(\lambda; x, y) = \frac{1}{\pi} \frac{\sin \left[ \sqrt{\lambda}(x - y) \right]}{x - y} \quad (9)$$

is the spectral function of the Fourier integral expansion, then the following uniform with respect to  $x \in \mathbb{R}$  estimate:

$$\|\theta(\lambda; x, \cdot) - \theta_0(\lambda; x, \cdot)\|_{L_q(\mathbb{R})} = O(1), \quad (10)$$

holds for sufficiently large values of  $\lambda$ .

In [8], we also proved that the spectral function  $\theta(\lambda; x, y)$  of the selfadjoint extension  $\mathcal{L}$  of the Hill operator (1) over  $\mathbb{R}$  satisfies the uniform with respect to  $(x, y) \in \mathbb{R} \times \mathbb{R}$  estimate

$$|\theta(\lambda; x, y) - \theta_0(\lambda; x, y)| = O(1). \quad (11)$$

Estimates of the spectral function both for ordinary differential operators and for partial differential operators were considered by many well-known mathematicians (Carleman, Gårding, Levitan, Hörmander, Kostyuchenko, Avakumovich, etc.). Usually, in order to estimate the spectral function of a differential operator these authors applied the method initially introduced by Carleman. It is based on a preliminary estimate of some function of operator and on the subsequent application of Tauberian theorems.

Such a method gives estimates of the spectral function only in the case when both arguments  $x, y$  vary on compact sets of the domain where the given operator is considered.

Development of the method introduced in the monograph [11] allowed us to estimate in [8] the spectral function  $\theta(\lambda; x, y)$  of the selfadjoint extension  $\mathcal{L}$  of the Hill operator over  $\mathbb{R}$  uniformly with respect to  $x, y$  varying independently over the infinite line  $\mathbb{R}$ .

Obtained estimates (10) and (11) of the spectral function give the necessary ground to prove [8], the uniform over the whole line  $\mathbb{R}$  equiconvergence of the spectral expansion of an arbitrary function  $f(x)$  from the class  $L_p(\mathbb{R})$  ( $1 \leq p \leq 2$ ), which corresponds to the selfadjoint extension  $\mathcal{L}$  of the Hill operator over  $\mathbb{R}$ , with the conventional Fourier integral expansion of the same function  $f(x)$ .

While using spectral expansions in various applications, it is important to know estimates for the deviation of the decomposed function  $f(x)$  from its spectral expansion  $\sigma_\lambda(x, f)$  as  $\lambda \rightarrow \infty$  (provided we know which class the function  $f(x)$  is from). Furthermore it is important to obtain these estimates in the uniform with respect to  $x \in \mathbb{R}$  metric.

This problem is covered in [9]. In that paper, we consider any bounded below selfadjoint extension  $\mathcal{L}$  of the Schrödinger operator

$$Hu = -u'' + q(x)u, \quad x \in \mathbb{R}, \quad (12)$$

with an arbitrary bounded and measurable over  $\mathbb{R}$  potential  $q(x)$ .

Some particular cases of this Schrödinger operator are of a special importance:

- (1) the operator (12) with a continuous periodic potential  $q(x)$  (the Hill operator),
- (2) the operator (12) with the potential  $q(x) = \beta \cos(\gamma x)$ , where  $\beta$  and  $\gamma$  are constants (the Mathieu-Hill operator),
- (3) the operator (12) with an almost periodic potential  $q(x)$ .

Without loss of generality, we assume this extension  $\mathcal{L}$  to be strictly positive with spectrum's infimum  $\lambda_0 \geq 1$ . Indeed, the general case of a selfadjoint bounded below extension  $\mathcal{L}$  can be reduced to it if we add a constant to the potential  $q(x)$  (after this transformation, the new potential is also bounded and measurable, while the whole spectrum shifts).

The following statement forms the main result of [9].

**MAIN THEOREM 2.** *Let  $\mathcal{L}$  be a selfadjoint and bounded below by  $\lambda_0 \geq 1$  extension of the Schrödinger operator (12) over  $\mathbb{R}$  with the potential  $q(x)$ , which is an arbitrary bounded and measurable over  $\mathbb{R}$  function. Let  $f(x)$  be any function from the Sobolev-Liouville class  $L_2^\alpha(\mathbb{R})$ , where  $\alpha \in (1/2, 2]$ , and  $\sigma_\lambda(x, f)$  be the spectral expansion of  $f(x)$  related to  $\mathcal{L}$ . Then the difference  $\sigma_\lambda(x, f) - f(x)$  satisfies the following uniform for  $x \in \mathbb{R}$  estimate:*

$$|\sigma_\lambda(x, f) - f(x)| = \lambda^{1/4-\alpha/2} \cdot o(1); \quad (13)$$

here  $o(1)$  denotes an entity that tends to zero as  $\lambda \rightarrow \infty$  uniformly over  $\mathbb{R}$ .

The estimate (13) clarifies the uniform over the whole line  $\mathbb{R}$  rate of vanishing of  $|\sigma_\lambda(x, f) - f(x)|$  as  $\lambda \rightarrow \infty$ .

The proof of Main Theorem 2 uses one auxiliary result that expresses the main characteristic property of the Fourier transforms of a function  $f(x)$  from the Sobolev-Liouville class  $L_2^\alpha(\mathbb{R})$ . This result also has a separate interest.

**THEOREM 3.** *Let  $\mathcal{L}$  be a selfadjoint bounded below by  $\lambda_0 \geq 1$  extension of the Schrödinger operator (12) over  $\mathbb{R}$  with the potential  $q(x)$ , which is an arbitrary bounded and measurable over  $\mathbb{R}$  function. Let  $f(x)$  be any function from the Sobolev-Liouville class  $L_2^\alpha(\mathbb{R})$  with a fixed derivation order  $\alpha \in (0, 2]$ . Then the Fourier transforms  $\hat{f}_i(t)$  of  $f(x)$  related to the ordered spectral representation<sup>4</sup> of  $L_2(\mathbb{R})$  space with respect to the extension  $\mathcal{L}$  satisfy the following inequality:*

$$\sum_{i=1}^m \int_{\lambda_0}^{\infty} |\hat{f}_i(t)|^2 \cdot t^\alpha dp(t) \leq C \|f\|_{L_2^\alpha(\mathbb{R})}^2. \quad (14)$$

Following the well-known Bernstein's Theorem on absolute and uniform convergence of the Fourier trigonometric series for a periodic function  $f(x)$  from the Hölder class  $C^\alpha[-\pi, \pi]$  with  $\alpha > 1/2$  (see, e.g., [12, Chapter 6, Theorem 3.1]), it is natural to introduce the notion of the absolute and uniform over the whole line  $\mathbb{R}$  convergence of the spectral expansion

$$\sigma_\lambda(x, f) = \sum_{i=1}^m \int_{\lambda_0}^{\lambda} \hat{f}_i(t) u_i(x, t) dp(t) \quad (15)$$

of an arbitrary function  $f(x)$ .

We say that the spectral expansion (15) converges absolutely and uniformly over the whole line  $\mathbb{R}$ , if the integral

$$\int_{\lambda_0}^{\infty} |\hat{f}_i(t)| \cdot |u_i(x, t)| dp(t)$$

converges for all  $x \in \mathbb{R}$  and  $i = \overline{1, m}$ , and there exists the uniform with respect to  $x \in \mathbb{R}$  limit

$$\lim_{\lambda \rightarrow \infty} \sum_{i=1}^m \int_{\lambda}^{\infty} |\hat{f}_i(t)| \cdot |u_i(x, t)| dp(t) = 0. \quad (16)$$

<sup>4</sup>This representation is characterized by the spectral measure  $p(t)$ , multiplicity  $m \leq 2$ , multiplicity sets  $e_i$ , and the generalized eigenfunctions  $u_i(x, t)$ .

**THEOREM 4 (GENERALIZED THEOREM OF BERNSTEIN'S TYPE).** *If  $\mathcal{L}$  is a selfadjoint bounded below by  $\lambda_0 \geq 1$  extension of the Schrödinger operator (12) over  $\mathbb{R}$  with a bounded and measurable over  $\mathbb{R}$  potential  $q(x)$ , then the corresponding spectral expansion (15) of an arbitrary function  $f(x)$  from the Sobolev-Liouville class  $L_2^\alpha(\mathbb{R})$  with  $\alpha > 1/2$  converges absolutely and uniformly over the whole line  $\mathbb{R}$ .*

Theorem 4 is also proved in [9]. Note that as in all other theorems of [8,9], the absolute and uniform convergence of the spectral expansion (15) is established *over the whole line  $\mathbb{R}$* .

In [9], we also prove that in the important case of the Mathieu-Hill operator

$$Hu = -u'' + \beta \cos(\gamma x)u, \quad x \in \mathbb{R}, \quad (17)$$

the restriction  $\alpha \leq 2$  in Main Theorem 2 and in Theorem 3 can be removed. For the Mathieu-Hill operator (17) the following statement holds.

**THEOREM 5.** *Let  $\mathcal{L}$  be a selfadjoint bounded below by  $\lambda_0 \geq 1$  extension of the Mathieu-Hill operator (17) over  $\mathbb{R}$ , and  $f(x)$  be any function from the Sobolev-Liouville class  $L_2^\alpha(\mathbb{R})$ . Then the estimate (14) for the Fourier transforms  $\hat{f}_i(t)$  of  $f(x)$  holds for any fixed  $\alpha \in (0, \infty)$ , and the uniform over the whole line  $\mathbb{R}$  estimate<sup>5</sup> (13) for the difference of the function  $f(x)$  and its spectral expansion  $\sigma_\lambda(x, f)$  holds for any fixed  $\alpha \in (1/2, \infty)$ .*

Thus in the important case of the Schrödinger operator, the Mathieu-Hill operator (17), one can get the desired rate of vanishing of  $|\sigma_\lambda(x, f) - f(x)|$  as  $\lambda \rightarrow \infty$  (uniformly with respect to  $x \in \mathbb{R}$ ), choosing the decomposed function  $f(x)$  from the Sobolev-Liouville class with a sufficiently high derivation order  $\alpha$ .

Concluding, we note that analysis of the proof of Main Theorem 1 (see [8]) and the estimate for the integral of squares of generalized eigenfunctions (see [13]) makes it possible to state that Main Theorem 1 holds true, also in the case when the Hill operator is substituted by the Schrödinger operator with a bounded and measurable potential  $q(x)$ .

## REFERENCES

1. E.A. Coddington and N. Levinson, *Theory of Ordinary Differential Equations*, McGraw-Hill, (1955).
2. V.A. Marchenko, *Sturm-Liouville Operators and Their Applications*, Naukova Dumka, Kiev, (1977).
3. V.A. Marchenko and I.V. Ostrovsky, (in Russian), *Matem. Sbornik* **97** (4), 540–606, (1975).
4. D. Schenk and M.A. Shubin, (in Russian), *Matem. Sbornik* **128** (4), 474–491, (1985).
5. M. Reed and B. Simon, *Methods of Modern Mathematical Physics, II*, Academic Press, New York, (1975).
6. M. Reed and B. Simon, *Methods of Modern Mathematical Physics, IV*, Academic Press, New York, (1978).
7. G.V. Rozenblyum, M.Z. Solomyak and M.A. Shubin, Spectral theory of differential operators, In *Itogi of Science and Technics: Modern Problems in Mathematics*, Volume 64, (in Russian), VINITI, Moscow, (1989).
8. V.A. Il'in and I. Antoniou, *Diff. Uravneniya* **31** (8), (1995).
9. V.A. Il'in and I. Antoniou, *Diff. Uravneniya*, (in preparation).
10. N. Dunford and J.T. Schwartz, *Linear Operators, II: Spectral Theory. Selfadjoint Operators in Hilbert Space*, Interscience, New York, (1957).
11. V.A. Il'in, *Spectral Theory of Differential Operators*, Nauka, Moscow, (1991).
12. A. Zygmund, *Trigonometric Series, I*, Cambridge, (1968).
13. V.A. Il'in and L.V. Kritskov, *Diff. Uravneniya* **31** (9), (1995).

<sup>5</sup>See Main Theorem 2 for the explanation of the notation  $o(1)$  in (13).